



Seacoast Orthopedics

Over 30 years of dedication to outstanding patient care

Hip Replacement Post-Op Discharge Instructions

Your Surgeon:

Andrew Banos, MD
Seacoast Orthopedics
21 Highland Ave, Suite 16
Newburyport MA 01950
Phone: (978) 462-7555
Website: <http://seacoastorthopedics.com/>



Seacoast Orthopedics

Over 30 years of dedication to outstanding patient care

Hip Replacement Post-Op Discharge Instructions

What to Expect:

It is normal to feel tired and washed out after surgery. Rest is important, but walking is also important to prevent problems and to regain your strength and energy. Pace yourself according to how you feel. Rest when you feel tired.

You may have a sore throat if you have had general anesthesia. This is normal and will go away in 1-2 days.

Your leg can remain painful and swollen for several weeks time, depending on the problem you have and the amount of surgery that was done. You may experience some bruising around your incision.

Pain Management:

You can expect to have pain for the first week or so after surgery. The pain is often worse at night. You will be given a prescription for pain medication upon discharge from the hospital. **This will be sent electronically to the pharmacy on file.** You should take this narcotic pain medication as directed for moderate to severe pain. You should take your pain medication with food. If you cannot tolerate this pain medication because of nausea or itching, please call the office. You should call the office for a refill several days before you run out of pain medication. **Refilling your new prescription may take 2-3 business days.** Narcotics will not be refilled by the on-call physician after hours or on weekends.

For mild pain, you may take acetaminophen (Tylenol). (However remember that medications such as Percocet, Vicodin, and Tylenol #3 contain acetaminophen and you should not take more than 3000 mg of acetaminophen daily.)

Pain medication can cause constipation. To make it easier to have a bowel movement you should drink extra fluids, eat foods high in fiber, or take a stool softener such as colace. Pain medications can also cause itching of the skin. For mild itching you can take Benadryl.

You should avoid non-steroidal anti-inflammatory medications like Motrin, Advil, Ibuprofen, Aleve or Naprosyn as these may cause bleeding.

Do not drive, drink alcohol or make important decisions while taking narcotic pain medication. Please call your doctor if the pain medication is not working.

Blood Clot (DVT) Prevention:

If you were given a prescription for Aspirin, you should take it twice daily for 4 weeks. You may obtain this medication over the counter at the pharmacy. (Do not take if you have a history of stomach ulcers, bleeding disorders or allergy to aspirin).

You were given TED compression stockings. It is important to continue wearing these stockings for the first two weeks after surgery. You should wear them at all times for as long as you can tolerate

them. However, you may remove the stockings as you become more mobile. You can remove them for showering and to wash them.

You should pump your ankles up and down several times an hour to keep the blood circulating in your leg. Don't lie in bed or on the couch for hours and hours at a time. It's good to move around (with the help of crutches or a walker) to get the blood flowing in your legs.

Treatment of Nausea:

If you received a Scopolamine patch (behind your ear) in the hospital, leave it in place for 24 hours. Be sure to wash your hands immediately after taking it off.

If you are nauseous or vomiting limit your intake to clear liquids only, and advance diet as tolerated. If still nauseous or vomiting after 12 hours, please call the office.

Dressing:

Keep your dressings clean and dry. Your dressings can be removed in 5 days. You may remove the bandages but **leave in place the underlying white steri-strips** that are covering the small incisions. The steri-strips will fall off on their own over time.

You may shower after 5 days, letting water gently wash over the incisions. Do not scrub the incisions. Do not submerge in a bath or pool until directed by your surgeon after the skin is fully healed. Antibiotic creams or other ointments should not be used.

Your sutures are typically under the skin and will dissolve on their own. You have surgical skin glue overlying the large incision. Do not scrub, pick, or peel off the glue. It will flake off over time. You may apply a new dressing over the incision for comfort if you wish, however you do not have to, as the tissue glue provides a protective barrier for the skin. You should keep the area clean either way.

Icing:

Ice bags can help to control swelling. Ice should be applied 30 minutes at a time every hour or two. Put a thin towel or T shirt on your skin if using ice in a plastic bag. Icing is most important in the first 72 hours, although many people find that continuing it lessens their postoperative pain weeks after surgery.

What Activities am I allowed to do at Home:

You should plan on taking a few weeks off from work. You should be up and walking as much as possible.

You should drink lots of fluids, and eat light foods at first like toast, crackers, soup and ginger ale.

Avoid smoking or caffeinated drinks as they impair healing.

You may not drive if you are taking pain medications. If you had a right hip replacement, you may not drive until you are walking normally without pain and can safely operate the brake pedal, this may take 4-6 weeks or longer.

Postoperative Visit:

You should have a follow up appointment scheduled for 2 weeks after surgery. If you do not, you should call the office the day after your surgery to schedule your post-operative visit.

When to Call:

If you have any trouble breathing or are having chest pain, you should dial 911 right away.

For any of the following danger signs, you should call your doctor's office: fever above 100.6 (after the first day or two – a slight fever can be normal the first day), redness or continued drainage from the incisions after 48hrs, persistent numbness, tingling, or discoloration of the foot, calf pain with swelling, or severe pain not relieved by pain medication.

Dental work after a joint replacement:


You should wait 3 months before having any dental procedures including routine cleaning due to the risk of infection of the joint replacement. After 3 months, dental procedures may be scheduled. Antibiotics are strongly recommended prior to any dental procedures. Your dentist may be able to order the antibiotic for you, if not please call our office to request a prescription at least one week prior to your dental appointment.

Hip Precautions:


It is very important to follow the precautions below while the tissues around your hip are healing to prevent dislocation of your hip replacement:

Posterior Hip Precautions


Do not bend forward past 90 degrees.



Do not cross your legs.



**Do not turn toes inward and
do not twist.**



**Do not turn knees
inward or together.**

