



## **Partial Knee Arthroplasty Post-Op Discharge Instructions**

### **Your Surgeon:**

Andrew Banos, MD  
Seacoast Orthopedics  
21 Highland Ave, Suite 16  
Newburyport MA 01950  
Phone: (978) 462-7555  
Website: <http://seacoastorthopedics.com/>



## Partial Knee Arthroplasty Post-Op Discharge Instructions

### **What to Expect:**

It is normal to feel tired and washed out after surgery. Rest is important, but walking is also important to prevent problems and to regain your strength and energy. Pace yourself according to how you feel. Rest when you feel tired.

You may have a sore throat if you have had general anesthesia. This is normal and will go away in 1-2 days.

If you have a nerve block, the local anesthetic may keep your leg numb for up to 24 hours.

Your leg can remain painful and swollen for several weeks time, depending on the problem you have and the amount of surgery that was done. You may experience some bruising around your incisions. You should elevate your knee higher than the level of your pelvis as often as possible to minimize the swelling.

### **Pain Management:**

You can expect to have pain for the first week or so after surgery. The pain is often worse at night. You will be given a prescription for pain medication upon discharge from the hospital. **This will be sent electronically to the pharmacy on file.** You should take this narcotic pain medication as directed for moderate to severe pain. You should take your pain medication with food. If you cannot tolerate this pain medication because of nausea or itching, please call the office. You should call the office for a refill several days before you run out of pain medication. **Refilling your new prescription may take 2-3 business days.** Narcotics will not be refilled by the on-call physician after hours or on weekends.

For mild pain, you may take acetaminophen (Tylenol). (However remember that medications such as Percocet, Vicodin, and Tylenol #3 contain acetaminophen and you should not take more than 3000 mg of acetaminophen daily.)

**Pain medication can cause constipation. To make it easier to have a bowel movement you should drink extra fluids, eat foods high in fiber, or take a stool softener such as colace.** Pain medications can also cause itching of the skin. For mild itching you can take Benadryl.

You should not take non-steroidal anti-inflammatory medication like Motrin, Advil, Ibuprofen, Aleve or Naprosyn as these may cause bleeding.

Do not drive, drink alcohol or make important decisions while taking narcotic pain medication. Please call your doctor if the pain medication is not working.

### **Treatment of Nausea:**

If you received a Scopolamine patch (behind your ear) in the hospital, leave it in place for 24 hours. Be sure to wash your hands immediately after taking it off.

If you are nauseous or vomiting limit your intake to clear liquids only, and advance diet as tolerated. If still nauseous or vomiting after 12 hours, please call the office.

### **Dressing:**

Keep your dressing clean and dry. Your dressing can be removed in 5 days. You should keep the incision clean, dry, and covered with a dressing for the first 2 weeks after surgery. You should change the dressing as needed, typically once daily.

Do not get the incision wet until seen at the 2 week follow up appointment. Do not submerge the incisions in water until directed by your surgeon after the skin is fully healed. Antibiotic creams or other ointments should not be used.

You should use 4"x4" or larger size gauze and a new ACE bandage for dressing changes. You may purchase these supplies at the drugstore. **Please do not use dressings with adhesive.** Repeated use of adhesive dressings around incisions on the knee can cause skin irritation which can increase the risk for infection.

You have staples underneath the dressing, which will be removed 10-14 days after surgery. Do not attempt to remove the staples on your own.

### **Icing:**

Apply ice bags or use the cryotherapy device you were given to control swelling. Ice should be applied 30 minutes at a time every hour or two. Put a thin towel or T shirt on your skin if using ice in a plastic bag. Icing is most important in the first 72 hours, although many people find that continuing it lessens their postoperative pain weeks after surgery.

### **Blood Clot (DVT) Prevention:**

If you were given a prescription for Aspirin, you should take it twice daily for 4 weeks. You may obtain this medication over the counter at the pharmacy. (Do not take if you have a history of stomach ulcers, bleeding disorders or allergy to aspirin).

You were given TED compression stockings. It is important to continue wearing these stockings for the first two weeks after surgery. You should wear them at all times for as long as you can tolerate them. However, you may remove the stockings as you become more mobile. You can remove them for showering and to wash them.

You should pump your ankles up and down several times an hour to keep the blood circulating in your leg. Don't lie in bed or on the couch for hours and hours at a time. It's good to move around (with the help of crutches) to get the blood flowing in your legs.

### **Exercises:**

Unless otherwise instructed by your surgeon you may put weight on the leg and bend the leg as tolerated. You should work with physical therapy on your knee range of motion.

You should not do any sports or athletics until cleared by your surgeon. You should pump your ankles up and down several times an hour to keep the blood circulating in your leg.

**What Activities am I allowed to do at Home:**

You should plan on taking some time off from work. You can resume work when the pain and swelling subside (this can be a several weeks or more depending on what type of work you do)

You should drink lots of fluids, and eat light foods at first like toast, crackers, soup and ginger ale.

Avoid smoking or caffeinated drinks as they impair healing.

**Postoperative Visit:**

You should have a follow up appointment scheduled. If you do not, you should call the office the day after your surgery to schedule your post-operative visit.

Plan to wear loose comfortable clothing your post-operative appointments with pant legs that easily roll above your knee so that the surgical incisions and knee can easily be examined. Alternatively, wear shorts underneath pants that can be easily removed so that the knee can be examined.

**When to Call:**

If you have any trouble breathing or are having chest pain, you should dial 911 right away.

For any of the following danger signs, you should call your doctor's office: fever above 100.6 (after the first day or two – a slight fever can be normal the first day), redness or drainage from the incision, numbness, tingling, or discoloration of the foot, if you have severe pain not relieved by pain medication, or if the numbness/tingling returns after the nerve block has worn off.

**Dental work after a joint replacement:**

You should wait 3 months before having any dental procedures including routine cleaning due to the risk of infection of the joint replacement. After 3 months, dental procedures may be scheduled. Antibiotics are strongly recommended prior to any dental procedures. Your dentist may be able to order the antibiotic for you, if not please call our office to request a prescription at least one week prior to your dental appointment.